



1. Consider the reaction



Suppose that the rate of loss of molecular nitrogen,  $-\frac{d[\text{N}_2]}{dt}$ , is 0.060 M/s at a particular time during the reaction. What is the rate of formation of ammonia,  $\frac{d[\text{NH}_3]}{dt}$ ?

- A. 0.020 M/s    B. 0.030 M/s    C. 0.060 M/s    **D. 0.12 M/s**    E. 0.18 M/s

2. For the overall chemical equation,



which one of the following can you rightly assume?

- A. The reaction is third-order overall.  
B. The reaction is second-order overall.  
C. The rate law is, rate =  $k[\text{H}_2\text{S}][\text{O}_2]$   
D. The rate law is, rate =  $k[\text{H}_2\text{S}]^2[\text{O}_2]$   
**E. The rate law cannot be determined from the information given.**

3. The reaction  $\text{A} + 2\text{B} \rightarrow \text{products}$  was found to have the rate law, rate =  $k[\text{A}][\text{B}]^2$ . Predict by what factor the rate of reaction will increase when the concentration of A is doubled and the concentration of B is also doubled.

- A. 2                      B. 4                      C. 6                      **D. 8**                      E. 9

4. Consider the hypothetical reaction  $\text{A} + 2\text{B} \rightarrow \text{products}$ . Use the following data to determine the **rate** of the reaction when  $[\text{A}] = 0.050 \text{ M}$  and  $[\text{B}] = 0.15 \text{ M}$ .

Expt. #	$[\text{A}]_0$	$[\text{B}]_0$	Initial rate
1	0.20	0.20	0.76 M/s
2	0.20	0.40	0.76 M/s
3	0.40	0.20	1.52 M/s

- A. 0.19 M/s**    B. 0.38 M/s    C. 0.57 M/s    D. 0.95 M/s    E. none of these

5. If concentration is expressed in units of moles per liter (mol/L) and time in units of seconds (s), the **units of a third-order rate constant** are

- A.  $\text{L mol}^{-1} \text{s}^{-1}$                       **B.  $\text{L}^2 \text{mol}^{-2} \text{s}^{-1}$**                       C.  $\text{s}^{-1}$   
D.  $\text{s}^{-2}$                                       E.  $\text{mol L}^{-1} \text{s}^{-1}$

6. A certain first-order reaction  $\text{A} \rightarrow \text{B}$  is 75% complete in 8.0 minutes at 25°C. What is the **half-life** of the reaction?

- A. 4.0 min**    B. 8.0 min    C. 16 min    D. 19 min    E. 39 min

7. For a certain reaction,  $A \rightarrow \text{products}$ , it was found that 20 minutes were required for the concentration of A to decrease from 1.0 M to 0.50 M, and that an additional 20 minutes were required for [A] to decrease from 0.50 M to 0.25 M. What is the **rate law** for this reaction?
- A.  $\text{rate} = k[A]^{\square 1}$   
 B.  $\text{rate} = k$   
 C.  $\text{rate} = k[A]$   
 D.  $\text{rate} = [A]^2$   
 E. cannot be determined from the information given

8. The rate constant for the second-order reaction



is  $0.54 \text{ M}^{-1}\text{s}^{-1}$  at  $300^\circ\text{C}$ . If the initial concentration of  $\text{NO}_2$  was 0.50 M, what would the **concentration of  $\text{NO}_2$**  be after 15 sec?

- A. **0.10 M**      B. 0.14 M      C. 0.21 M      D. 0.28 M      E. 0.39 M

9. A certain zero-order reaction  $A \rightarrow B$  for which  $[A]_0 = 1.0 \text{ M}$  is 30% complete in 13 seconds at  $25^\circ\text{C}$ . What the **rate constant, k**, for this reaction at  $25^\circ\text{C}$ ?

- A. 0.012 M/s    B. **0.023 M/s**    C. 0.035 M/s    D. 0.046 M/s    E. 0.058 M/s

10. The isomerization of cyclopropane follows first-order kinetics. The rate constant at 600 K is  $2.72 \times 10^{-7} \text{ min}^{-1}$ , and the activation energy for the reaction is 270 kJ/mol. Calculate the value of the **rate constant** (in  $\text{min}^{-1}$ ) at 700 K.

- A.  **$6.20 \times 10^{14}$**                       B.  $2.05 \times 10^{11}$                       C.  $1.86 \times 10^1$   
 D.  $6.87 \times 10^2$                       E.  $1.32 \times 10^4$

11. For a certain second-order reaction, rate constant (k) at  $25^\circ\text{C}$  is  $0.235 \text{ M}^{-1} \text{ s}^{-1}$ . At  $50^\circ\text{C}$  the rate constant is  $1.12 \text{ M}^{-1}\text{s}^{-1}$ . What is the **frequency factor, A**, for this reaction in units of  $\text{M}^{-1}\text{s}^{-1}$ ?

- A.  $4.26 \square 10^4$     B.  $3.21 \square 10^5$     C.  $2.41 \square 10^6$     D.  $1.82 \square 10^7$     E.  **$1.37 \square 10^8$**

12. If a catalyst could be found that would lower the activation energy by 25.0 kJ/ mol for a particular reaction, by what **factor** would the rate constant for this reaction be increased at  $25^\circ\text{C}$ ? (Assume the frequency factor remains the same.)

- A.  $5.66 \times 10^1$     B.  $4.26 \times 10^2$     C.  $3.20 \times 10^3$     D.  **$2.41 \times 10^4$**     E.  $1.81 \times 10^5$