

**MATH 285 E1/F1 GRADED HOMEWORK SET 4**  
**DUE FRIDAY OCTOBER 24 IN LECTURE**

This time, the homework has **just one part**. Please staple your homework together, and put your **name and section** on it. *Thank you!*

- (1) (10 points) Find the *general solution* of the differential equation

$$(D - 3)y = xe^{2x} + e^{3x}$$

*Hint:* Use the annihilator method.

- (2) (5 points) Consider the forced mass-spring oscillator with mass  $m = 4$ , no damping  $c = 0$ , and a spring constant  $k$  that is adjustable. Consider a driving forces that is the sum of two cosine functions

$$F(t) = \cos 2t + \cos 5t.$$

The differential equation for the displacement  $x(t)$  is then

$$4x'' + kx = \cos 2t + \cos 5t.$$

For what values of  $k$  is it impossible to find a particular solution of the form  $A \cos 2t + B \cos 5t$ ? (These are the values of  $k$  for which resonance occurs.)

- (3) (5 points) Recall from the lecture the damped ( $c > 0$ ) forced oscillator obeying the differential equation

$$mx'' + cx' + kx = F_0 \cos \omega t.$$

In the lecture, we derived a particular solution of this equation

$$x_p(t) = C \cos(\omega t - \alpha),$$

where

$$C = \frac{F_0}{\sqrt{(k - m\omega^2)^2 + (c\omega)^2}}, \quad \tan \alpha = \frac{c\omega}{k - m\omega^2}.$$

Here is something we did not talk about in the lecture: the solution  $x_p(t)$  is called the *steady-state* solution, because any solution will asymptotically converge to it as  $t \rightarrow \infty$ . More precisely, if  $x_1(t)$  is *any solution* of the same nonhomogeneous equation, then

$$\lim_{t \rightarrow \infty} [x_1(t) - x_p(t)] = 0.$$

Explain why this is true. You should use the description of the solutions of the homogeneous equation  $mx'' + cx' + kx = 0$  on page 191 of the textbook. *Hint:* It is very important that  $c > 0$ .

- (4) (5 points) Use the orthogonality properties of sine and cosine to compute the following integral

$$\int_{-\pi}^{\pi} (5 \sin 2t - \cos 3t)(\cos 2t + 5 \sin 6t + \sin 2t) dt$$