



System (4) has a unique solution provided the **determinant of coefficients**  $\Delta = \det(A)$  is nonzero, in which case the solution is given by

$$(5) \quad x_1 = \frac{\Delta_1}{\Delta}, \quad x_2 = \frac{\Delta_2}{\Delta}, \quad \dots, \quad x_n = \frac{\Delta_n}{\Delta}.$$

The determinant  $\Delta_j$  equals  $\det(B_j)$  where matrix  $B_j$  is matrix  $A$  with column  $j$  replaced by  $\vec{b} = (b_1, \dots, b_n)$ , which is the right side of system (4). The result is called **Cramer's Rule** for  $n \times n$  systems. Determinants will be defined shortly; intuition from the  $2 \times 2$  case and Sarrus' rule should suffice for the moment.

**Determinant Notation for Cramer's Rule.** The **determinant of coefficients** for system  $A\vec{x} = \vec{b}$  is denoted by

$$(6) \quad \Delta = \begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{vmatrix}.$$

The other  $n$  determinants in Cramer's rule (5) are given by

$$(7) \quad \Delta_1 = \begin{vmatrix} b_1 & a_{12} & \cdots & a_{1n} \\ b_2 & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ b_n & a_{n2} & \cdots & a_{nn} \end{vmatrix}, \quad \dots, \quad \Delta_n = \begin{vmatrix} a_{11} & a_{12} & \cdots & b_1 \\ a_{21} & a_{22} & \cdots & b_2 \\ \vdots & \vdots & \cdots & \vdots \\ a_{n1} & a_{n2} & \cdots & b_n \end{vmatrix}.$$

The literature is filled with conflicting notations for matrices, vectors and determinants. The reader should take care to use vertical bars *only* for determinants and absolute values, e.g.,  $|A|$  makes sense for a matrix  $A$  or a constant  $A$ . For clarity, the notation  $\det(A)$  is preferred, when  $A$  is a matrix. The notation  $|A|$  implies that *a determinant is a number*, computed by  $|A| = \pm A$  when  $n = 1$ , and  $|A| = a_{11}a_{22} - a_{12}a_{21}$  when  $n = 2$ . For  $n \geq 3$ ,  $|A|$  is computed by similar but increasingly complicated formulas; see Sarrus' rule and the *four properties* below.

**Sarrus' Rule for  $3 \times 3$  Matrices.** College algebra supplies the following formula for the determinant of a  $3 \times 3$  matrix  $A$ :

$$(8) \quad \begin{aligned} \det(A) &= \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \\ &= a_{11}a_{22}a_{33} + a_{21}a_{32}a_{13} + a_{31}a_{12}a_{23} \\ &\quad - a_{11}a_{32}a_{23} - a_{21}a_{12}a_{33} - a_{31}a_{22}a_{13}. \end{aligned}$$

The number  $\det(A)$  can be computed by an algorithm similar to the one for  $2 \times 2$  matrices, as in Figure 10. We remark that no further generalizations are possible: *there is no Sarrus' rule for  $4 \times 4$  or larger matrices!*




Figure 10. Sarrus' rule for  $3 \times 3$  matrices, which gives

$$\det(A) = (a + b + c) - (d + e + f).$$

**College Algebra Definition of Determinant.** The impractical definition is the formula

$$(9) \quad \det(A) = \sum_{\sigma \in S_n} (-1)^{\text{parity}(\sigma)} a_{1\sigma_1} \cdots a_{n\sigma_n}.$$

In the formula,  $a_{ij}$  denotes the element in row  $i$  and column  $j$  of the matrix  $A$ . The symbol  $\sigma = (\sigma_1, \dots, \sigma_n)$  stands for a rearrangement of the subscripts  $1, 2, \dots, n$  and  $S_n$  is the set of all possible rearrangements. The nonnegative integer  $\text{parity}(\sigma)$  is determined by counting the minimum number of pairwise interchanges required to assemble the list of integers  $\sigma_1, \dots, \sigma_n$  into natural order  $1, \dots, n$ .

Formula (9) reproduces the definition for  $3 \times 3$  matrices given in equation (8). We will have no computational use for (9). For computing the value of a determinant, see below *four properties* and *cofactor expansion*.

**Four Properties.** The definition of determinant (9) implies the following four properties:

<b>Triangular</b>	The value of $\det(A)$ for either an upper triangular or a lower triangular matrix $A$ is the product of the diagonal elements: $\det(A) = a_{11}a_{22} \cdots a_{nn}$ .
<b>Swap</b>	If $B$ results from $A$ by swapping two rows, then $\det(A) = (-1)\det(B)$ .
<b>Combination</b>	The value of $\det(A)$ is unchanged by adding a multiple of a row to a different row.
<b>Multiply</b>	If one row of $A$ is multiplied by constant $c$ to create matrix $B$ , then $\det(B) = c\det(A)$ .

It is known that these four rules suffice to compute the value of any  $n \times n$  determinant. The proof of the four properties is delayed until page 320.