
Indefinite Integrals, Substitution

We first lay out the example of the *indefinite integral*.

Let us consider the integral

$$\int_0^4 e^x dx.$$

From the Fundamental Theorem of Calculus, we know that the antiderivative of e^x is itself, so

$$\int_0^4 e^x dx = e^x \Big|_{x=0}^{x=4} = e^4 - e^0 = e^4 - 1.$$

But we also have

$$\int_2^3 e^x dx = e^x \Big|_{x=2}^{x=3} = e^3 - e^2,$$

and

$$\int_{-2}^7 e^x dx = e^x \Big|_{x=-2}^{x=7} = e^7 - e^{-2}.$$

And we see that in all these cases, the formula into which we are plugging in is the same; we just use different values to plug in. Moreover, as we will see below, it is typical that the vast majority of the work that we need to do will involve finding this function, and the plugging in of the boundary values is the easy part.

As such, we would like to just refer as this antiderivative function, and when we need to do so we will talk about the indefinite integral, which is just another word for antiderivative.

The one thing to point out here is that an antiderivative is not unique. For example, we know that

$$\frac{d}{dx}(x^3) = \frac{d}{dx}(x^3 + 2) = \frac{d}{dx}(x^3 - 12) = \frac{d}{dx}(x^3 + \pi^2 e^{17} \cos \sin(4)) = 3x^2.$$

In particular, this means that we can always add any constant we like to an antiderivative, since all of the functions

$$x^2, \quad x^3 + 2, \quad x^3 - 12, \quad x^3 + \pi^2 e^{17} \cos \sin(4)$$

are all antiderivatives of $3x^2$. In general, we will write this as

$$x^3 + C,$$

where C is an arbitrary constant.

And, again, we stress that the choice of constant C will not matter when computing a definite integral, since

$$\int_{-2}^7 e^x dx = e^x + C \Big|_{x=-2}^{x=7} = (e^7 + C) - (e^{-2} + C) = e^7 - e^{-2} + (C - C) = e^7 - e^{-2}.$$

So, the hard part is finding antiderivatives. Basically, all of the rules we come up with are just rules for differentiation run backwards! Examples:

1.
$$\int x^n = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1$$
2.
$$\int e^x = e^x + C$$
3.
$$\int \frac{1}{x} = \ln |x| + C$$
4.
$$\int cf(x) dx = c \int f(x),$$
5.
$$\int (f(x) + g(x)) dx = \int f(x) + \int g(x),$$

Unfortunately, there isn't a rule like

$$\int f(x)g(x) dx \neq \int f(x) dx \int g(x) dx.$$

There is a way to use the product rule in reverse, which we will see below — it is called “integration by parts”.

However, we can use the Chain Rule in reverse pretty easily. Say that $F'(x) = f(x)$, and then

$$\frac{d}{dx} F(g(x)) = F'(g(x)) \cdot g'(x) = f(g(x)) \cdot g'(x).$$

This means that

$$\int f(g(x)) \cdot g'(x) dx = F(g(x)) + C.$$

This gives rise to the technique of ***u*-substitution**.

Example 27. We want to compute

$$\int x\sqrt{3+x^2} dx.$$

Let us try $u = 3 + x^2$, which gives $du = 2x dx$, so that $x dx = du/2$. We then have

$$\int x\sqrt{3+x^2} dx = \int \frac{1}{2}\sqrt{u} du = \frac{2}{3}u^{3/2} + C = \frac{2}{3}(3+x^2)^{3/2} + C.$$

A crucial feature that made this work is that there was an extra power of x in the integrand. For example, if we were given

$$\int \sqrt{3+x^2} dx,$$

and tried this trick, we would get

$$\int \sqrt{3+x^2} dx = \int \sqrt{u} \frac{du}{2x},$$

and now what do we do?

Example 28. Let's try the integral

$$\int e^{x+e^x} dx.$$

Now, we might first think that our sub should be $u = x + e^x$, which gives $du = (1 + e^x) dx$, so we have

$$\int e^{x+e^x} dx = \int e^u \frac{dx}{1 + e^x},$$

which doesn't seem that helpful. But, let us go back to the original integral and multiply out, recalling that $e^{a+b} = e^a e^b$, so we have

$$\int e^{x+e^x} dx = \int e^x \cdot e^{e^x} dx,$$

and now we might see that it makes sense to use $u = e^x$, $du = e^x dx$, and then we have

$$\int e^{x+e^x} dx = \int e^u du = e^u + C = e^{e^x} + C.$$

Example 29. Sometimes we might have to do a lot more work than we originally thought! Let's try

$$\int \frac{x}{1 + \sqrt{1 - x^2} - x^2} dx.$$

Let us try $u = 1 - x^2$, so $du = -2x dx$, and we have

$$\int \frac{x}{1 + \sqrt{1 - x^2} - x^2} dx = -\frac{1}{2} \int \frac{du}{u + \sqrt{u}}.$$

Since we have a \sqrt{u} term, let us choose $v = \sqrt{u}$, and thus $dv = 1/(2\sqrt{u}) du$, or $du = 2v dv$, and we have

$$\int \frac{x}{1 + \sqrt{1 - x^2} - x^2} dx = - \int \frac{v dv}{v^2 + v} = - \int \frac{dv}{v + 1}.$$

Finally, choose $w = v + 1$ and $dw = dv$, and we have

$$\int \frac{x}{1 + \sqrt{1 - x^2} - x^2} dx = - \int \frac{dw}{w} = - \ln |w| + C.$$

Unpacking all of the definitions, we have

$$\begin{aligned} \int \frac{x}{1 + \sqrt{1 - x^2} - x^2} dx &= - \ln |w| + C \\ &= - \ln |v + 1| + C \\ &= - \ln |\sqrt{u} + 1| + C \\ &= - \ln |\sqrt{1 - x^2} + 1| + C. \end{aligned}$$

Also, if we are doing definite integrals, then we can also use integration by parts, but we should change the limits of integration as we go.