

Problem Set #7
 Chapter 10, Ex. 1,3,4, p.172;
 Chapter 11: Ex. 1,2,3,4, p.177

November 2, 2004

Ex. 1, p. 172

$$\frac{d}{dt} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x - x^2 - 2xy \\ 2y - y^2 - 3xy \end{pmatrix}$$

Equilibrium points are: $\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 2 \end{pmatrix}, \begin{pmatrix} 0.6 \\ 0.2 \end{pmatrix}$.

$$\mathcal{D} = \begin{pmatrix} 1 - 2x - 2y & -2x \\ -3y & 2 - 2y - 3x \end{pmatrix}$$

Equilib.point	Matrix	$Det(\mathcal{D})$	$Tr(\mathcal{D})$	Stability
$\begin{pmatrix} 0 \\ 0 \end{pmatrix}$	$\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	2 (2 > 0)	3 (3 > 0)	Unstable
$\begin{pmatrix} 0 \\ 2 \end{pmatrix}$	$\begin{pmatrix} -3 & 0 \\ -6 & -2 \end{pmatrix}$	6 (6 > 0)	-5 (-5 < 0)	Stable
$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$	$\begin{pmatrix} -1 & -2 \\ 0 & -1 \end{pmatrix}$	1 (1 > 0)	-2 (-2 < 0)	Stable
$\begin{pmatrix} 0.6 \\ 0.2 \end{pmatrix}$	$\begin{pmatrix} -0.6 & -1.2 \\ -0.6 & -0.2 \end{pmatrix}$	-0.6 (-0.6 < 0)	-0.8 (-0.8 < 0)	Unstable

Ex. 3, p.172

- $h = x^2y^3$
 - $\frac{\partial h}{\partial x} = 2y^3x$
 - $\frac{\partial h}{\partial y} = 3x^2y^2$
 - $\frac{\partial^2 h}{\partial x^2} = 2y^3$
 - $\frac{\partial^2 h}{\partial y^2} = 6x^2y$
 - $\frac{\partial^2 h}{\partial y \partial x} = 6xy^2$

$$\frac{\partial^2 h}{\partial x \partial y} = 6xy^2$$

The last two are equal.

- $h = x \cos(xy)$

$$\frac{\partial h}{\partial x} = -xy \sin(xy) + \cos(xy)$$

$$\frac{\partial h}{\partial y} = -x^2 \sin(xy)$$

$$\frac{\partial^2 h}{\partial x^2} = -xy^2 \cos(xy) - 2y \sin(xy)$$

$$\frac{\partial^2 h}{\partial y^2} = -x^3 \cos(xy)$$

$$\frac{\partial^2 h}{\partial y \partial x} = -2x \sin(xy) - x^2 y \cos(xy)$$

$$\frac{\partial^2 h}{\partial x \partial y} = -2x \sin(xy) - x^2 y \cos(xy)$$

The last two are equal.

- $h = \sin(x + y^2)$

$$\frac{\partial h}{\partial x} = \cos(x + y^2)$$

$$\frac{\partial h}{\partial y} = 2y \cos(x + y^2)$$

$$\frac{\partial^2 h}{\partial x^2} = -\sin(x + y^2)$$

$$\frac{\partial^2 h}{\partial y^2} = 2\cos(x + y^2) - 4y^2 \sin(x + y^2)$$

$$\frac{\partial^2 h}{\partial y \partial x} = -2y \sin(x + y^2)$$

$$\frac{\partial^2 h}{\partial x \partial y} = -2y \sin(x + y^2)$$

The last two are equal.

- $h = xe^y$

$$\frac{\partial h}{\partial x} = e^y$$

$$\frac{\partial h}{\partial y} = xe^y$$

$$\frac{\partial^2 h}{\partial x^2} = 0$$

$$\frac{\partial^2 h}{\partial y^2} = xe^y$$

$$\frac{\partial^2 h}{\partial y \partial x} = e^y$$

$$\frac{\partial^2 h}{\partial x \partial y} = e^y$$

The last two are equal.

Thus, we have seen through these examples that $\frac{\partial^2 h}{\partial y \partial x} = \frac{\partial^2 h}{\partial x \partial y}$.

Ex. 4, p. 172

Integrate $h(x, y)$ over the indicated rectangle.

a)

$$\int_{-1}^2 \int_0^1 dx dy = \int_{-1}^2 x|_0^1 dy = y|_{-1}^2 = 3$$

b)

$$\int_0^1 \int_{-1}^1 x dx dy = \int_0^1 (x^2/2)|_{-1}^1 dy = 0$$

c)

$$\int_0^1 \int_0^1 (x+y) dx dy = \int_0^1 (x^2/2 + yx)|_0^1 dy = \int_0^1 (1/2 + y) dy = (y/2 + y^2/2)|_0^1 = 1/2 + 1/2 = 1$$

d)

$$\int_2^3 \int_{-2}^1 xy dx dy = \int_2^3 (x^2 y/2)|_{-2}^1 dy = \int_2^3 \frac{y}{2}(1-4) dy = \int_2^3 -\frac{3}{2}y dy = (-3y^2/4)|_2^3 = -15/4$$

e)

$$\int_0^1 \int_0^1 \cos(xy) dx dy = \int_0^1 \frac{\sin(xy)}{y} \Big|_0^1 dy = \int_0^1 \frac{\sin y}{y} dy = 0.946$$

Chapter 11. Ex. 1, p.177

a) $M = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$

$$\vec{v} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$M \vec{v} = \begin{pmatrix} 1 \cdot 1 + 2 \cdot 1 \\ 0 \cdot 1 + 1 \cdot 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$$

b) $M = \begin{pmatrix} 3 & 2 \\ -5 & 4 \end{pmatrix}$

$$\vec{v} = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$$

$$M \vec{v} = \begin{pmatrix} 2 \\ 7 \end{pmatrix}$$

c) $M = \begin{pmatrix} 0.5 & 0.2 \\ 8 & 0.4 \end{pmatrix}$

$$\vec{v} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$M \vec{v} = \begin{pmatrix} 0.2 \\ 0.4 \end{pmatrix}$$