

MATH 21b Practice Questions

Problem 1.

Circle T if the given assertion is true, and circle F if it is false. There is no need to justify your answer.

- T F a) Let $A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Then $AB = BA$.
- T F b) Suppose \vec{u} is a non-zero vector in \mathbb{R}^n . The map of \mathbb{R}^n to itself that sends any given vector \vec{v} to $T(\vec{v}) = \vec{v} + \vec{u}$ is a linear transformation.
- T F c) For any square matrix A , the kernel of A^2 is a subspace of the kernel of A^3 .
- T F d) If $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $S: \mathbb{R}^m \rightarrow \mathbb{R}^n$ are linear transformations such that the kernel of S has dimension greater than zero, then the kernel of ST must have dimension greater than zero.
- T F e) If a system of linear equations has more unknowns than equations, there are always an infinite number of solutions.
- T F f) If the product of two matrices is 0, then one or the other must also be 0.
- T F g) If A is a matrix, then $\ker(A)$ must be the same subspace as $\ker(\text{rref}(A))$.
- T F h) If A is a matrix, then $\text{image}(A)$ must be the same as $\text{image}(\text{rref}(A))$.
- T F i) If A is a square matrix with linearly independent columns, then the rows are also linearly independent vectors.
- T F j) A linear transformation of \mathbb{R}^2 cannot send $\begin{pmatrix} 6 \\ -9 \end{pmatrix}$ to $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$ and $\begin{pmatrix} -2 \\ 3 \end{pmatrix}$ to $\begin{pmatrix} -2 \\ 1 \end{pmatrix}$.

Problem 2.

Let T denote a linear transformation of \mathbb{R}^2 that sends $\begin{pmatrix} 1 \\ -3 \end{pmatrix}$ to $\begin{pmatrix} 1 \\ -2 \end{pmatrix}$ and $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$ to $\begin{pmatrix} 0 \\ -1 \end{pmatrix}$.

- a) What are the coordinates of $\begin{pmatrix} 0 \\ -1 \end{pmatrix}$ with respect to the basis $\vec{v}_1 = \begin{pmatrix} 1 \\ -3 \end{pmatrix}$ and $\vec{v}_2 = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$.
- b) Write down the matrix of T with respect to the basis $\vec{v}_1 = \begin{pmatrix} 1 \\ -3 \end{pmatrix}$ and $\vec{v}_2 = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$.
- c) Write down the matrix of T with respect to the standard basis, $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

Problem 3.

Let A denote the matrix $\begin{pmatrix} 1 & 2 & 0 & 2 & 5 \\ 1 & 2 & 1 & -1 & 0 \\ 2 & 4 & -1 & 1 & 3 \\ 3 & 6 & -1 & -1 & 0 \end{pmatrix}$.

- Compute $\text{ref}(A)$.
- Give a basis for $\text{kernel}(A)$.
- Give a basis for $\text{image}(A)$.
- What is the dimension of $\text{kernel}(A^T)$?

Problem 4.

Let A denote the matrix $\begin{pmatrix} 1 & 2 \\ 0 & 1 \\ -1 & 0 \end{pmatrix}$.

- Give an orthonormal basis for the image of A .
- Give the matrix T (with respect to the standard basis of \mathbb{R}^3) that represents the orthogonal projection onto the image of A .
- Let \vec{v}_1, \vec{v}_2 denote any orthonormal basis for $\text{image}(A)$, and let \vec{v}_3 denote a unit length vector that is orthogonal to $\text{image}(A)$. Find the matrix with respect to the basis $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ for the orthogonal projection to $\text{image}(A)$.
- Write a non-zero vector that is orthogonal to $\text{image}(A)$.
- Write down the matrix (with respect to the standard basis of \mathbb{R}^3) that represents the orthogonal projection onto the orthogonal complement of $\text{image}(A)$.

Problem 5.

Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ denote the linear transformation whose matrix with respect to the

standard basis of \mathbb{R}^3 is $A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 2 \\ 3 & 1 & 2 \end{pmatrix}$. Meanwhile, let $\vec{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$, $\vec{v}_2 = \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}$, $\vec{v}_3 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$.

- Compute A^2 .
- Prove that \vec{v}_1, \vec{v}_2 , and \vec{v}_3 are a basis for \mathbb{R}^3 .
- Write down the vector $A\vec{v}_1$.

Answers

1. a) F b) F c) T d) F e) F f) F g) T h) F i) T j) T.

2. a) $\vec{v}_1 + \vec{v}_2$ b) $\begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}$ c) $\begin{pmatrix} -2 & -1 \\ 7 & 3 \end{pmatrix}$.

3. a) $\begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$ b) $\begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \\ -1 \\ -2 \\ 1 \end{pmatrix}$ c) The last 3 columns of A span the image

d) Since the kernel of A^T is the orthogonal complement to the image of A, and the image of A has dimension 3, the kernel of A^T has dimension 2.

4. a) $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$ and $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$.

b) $T = \frac{1}{8} \begin{pmatrix} 5 & 2 & -1 \\ 2 & 2 & 2 \\ -1 & 2 & 5 \end{pmatrix}$.

c) The matrix is $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$.

d) $\vec{v} = \frac{1}{\sqrt{5}} \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$.

e) The matrix is $\frac{1}{8} \begin{pmatrix} 1 & -2 & 1 \\ -2 & 4 & -2 \\ 1 & -2 & 1 \end{pmatrix}$