

**Estimation of the Second-order Intensities of a
Bivariate Stationary Point Process**

By DAVID R. BRILLINGER

The University of California, Berkeley

[Received October 1974. Final revision May 1975]

SUMMARY

We consider histogram and smoothed histogram type estimates of the auto and cross intensity functions of a bivariate stationary point process. The asymptotic distributions are found to be multiples of Poissons in the histogram case and linear combinations of Poissons in the smoothed case. These asymptotic distributions suggest the plotting of the square roots of the estimates in order to stabilize the variance and to make the distributions more nearly normal. Two examples of such plots are presented in the paper.

Keywords: AUTOINTENSITY FUNCTION; CROSSINTENSITY FUNCTION; POINT PROCESS; SQUARE ROOT TRANSFORMATION; STATIONARY PROCESS

1. INTRODUCTION

LET $\{N_1(t), N_2(t)\}$, $-\infty < t < \infty$, be a bivariate stationary point process with $N_1(t)$ being the number of events of Type 1 that occurred in the time interval $(0, t]$ and $N_2(t)$ the number of events of Type 2 in the same interval. Suppose that the process is orderly in the sense that there is zero probability that events occur simultaneously. The intensity of events of type a is defined by

$$p_a = \lim_{h \downarrow 0} \Pr \{\text{type } a \text{ event in } (t, t+h]\} / h \quad (1.1)$$

for $a = 1, 2$. The existence of the limit (1.1) was shown by Khintchine (1960). Korolyuk showed that, with orderliness,

$$E\{dN_a(t)\} = p_a dt \quad (1.2)$$

(see Khintchine, 1960). The second-order product density function of events of type a with events of type b is defined by

$$p_{ab}(u) = \lim_{h, h' \downarrow 0} \Pr \{\text{type } a \text{ event in } (t+u, t+u+h] \text{ and type } b \text{ event in } (t, t+h']\} / (hh') \quad (1.3)$$

for $a, b = 1, 2$ and $u \neq 0$. The second-order intensity function of events of type a , given events of type b , is defined by

$$\begin{aligned} m_{ab}(u) &= \lim_{h \downarrow 0} \Pr \{\text{type } a \text{ event in } (t+u, t+u+h] \mid \text{type } b \text{ at } t\} / h \\ &= p_{ab}(u) / p_b \end{aligned} \quad (1.4)$$

for $a, b = 1, 2$ and $u \neq 0$. In this paper we are concerned with large sample properties of estimates of $p_{ab}(u)$, $m_{ab}(u)$ that have the form proposed in Griffith and Horn (1963), Cox (1965) and Cox and Lewis (1972). We shall propose a modified form of these estimates and, in the light of the large sample properties, recommend the application of a square root transformation. Numerous practical examples of estimates of the original form may be found in Bryant *et al.* (1973) for bivariate processes consisting of the input and output spike trains of nerve cells. Two examples of the modified estimates are presented in this paper. Numerous additional examples are given in a paper by Brillinger, Bryant and Segundo which is in preparation.

Suppose that the process $\{N_1(t), N_2(t)\}$ is given for $0 < t \leq T$, that is, the times at which events occurred in the interval $(0, T]$ are known. Let the times of events of type a be s_1, s_2, \dots , and the times of events of type b be t_1, t_2, \dots . Let $\beta > 0$ denote a scale parameter. Next, let $\# \{A\}$ denote the number of elements in a set A . Then the estimates of $p_{ab}(u)$ and $m_{ab}(u)$, considered in Cox and Lewis (1972), are based on the counting variate

$$J_{ab}^T(u) = \# \{(j, k) \text{ such that } u - \beta < s_j - t_k < u + \beta \text{ and } s_j \neq t_k\}. \tag{1.5}$$

$J_{ab}^T(u)$ counts the number of a events falling in a cell of bin width 2β and midpoint u time units along from a b event. It is a histogram type statistic. Cox and Lewis (1972) show that

$$\begin{aligned} EJ_{ab}^T(u) &\doteq (T - u) \int_{u - \beta}^{u + \beta} p_{ab}(v) dv \\ &\doteq 2\beta T p_{ab}(u) \end{aligned} \tag{1.6}$$

for large T , small β and moderate u , suggesting the estimates

$$\left. \begin{aligned} \hat{p}_{ab}(u) &= J_{ab}^T(u) / (2\beta T), \\ \hat{m}_{ab}(u) &= J_{ab}^T(u) / \{2\beta N_b(T)\}. \end{aligned} \right\} \tag{1.7}$$

We shall determine the asymptotic distributions of these estimates under certain regularity conditions. In addition we shall propose the use of the following modified estimates

$$\left. \begin{aligned} \hat{p}'_{ab}(u) &= \hat{p}_{ab}(u) + |u| N_a(T) N_b(T) / T^3, \\ \hat{m}'_{ab}(u) &= \hat{m}_{ab}(u) + |u| N_a(T) / T^2 \end{aligned} \right\} \tag{1.8}$$

for $|u| \leq T$. Under the regularity conditions mentioned, these appear to have better overall mean-squared error properties. Their definition will be motivated in Section 3. In the case that $|u|$ is not large compared to T , there is little difference between the estimates of (1.7) and (1.8). Their asymptotic distributions are the same.

2. THE ASYMPTOTIC DISTRIBUTIONS

Many random processes that occur in practice seem to satisfy some form of mixing condition, that is, functionals of the process that are well separated in time are only weakly dependent. We will make use of the following condition of that character.

Definition. A stationary bivariate process $\{N_1(t), N_2(t)\}$, $-\infty < t < \infty$, is called strong mixing when

$$\alpha(\tau) = \sup \{ |P(AB) - P(A)P(B)| : A \in \mathcal{M}_{-\infty}^t, B \in \mathcal{M}_{t+\tau}^\infty \} \rightarrow 0 \tag{2.1}$$

as $\tau \rightarrow \infty$. Here $P(\cdot)$ denotes the probability measure of the process and \mathcal{M}_u^v denotes the σ -algebra of events generated by events of the form

$$\{N_{a_1}(v_1) - N_{a_1}(u_1) \leq h_1, \dots, N_{a_K}(v_K) - N_{a_K}(u_K) \leq h_K\},$$

where $a_k = 1, 2$; $u < u_k < v_k \leq v$; h_k is a non-negative integer for $k = 1, 2, \dots, K$ and $K = 1, 2, \dots$

This condition appears in Volkonskii and Rozanov (1959) for example. We shall also require that the second- to fourth-order moments of the process have the following forms:

$$\left. \begin{aligned} E\{dN_a(t+u) dN_b(t)\} &= p_{ab}(u) dt du, \\ E\{dN_a(t+u) dN_b(t+v) dN_c(t)\} &= p_{abc}(u, v) dt du dv, \\ E\{dN_a(t+u) dN_b(t+v) dN_c(t+w) dN_d(t)\} &= p_{abcd}(u, v, w) dt du dv dw \end{aligned} \right\} \tag{2.2}$$

for $a, b, c, d = 1, 2$ and $u, v, w, 0$ distinct. Finally, let $P(\mu)$ denote a Poisson variate with mean μ .

Theorem 1. Let $\{N_1(t), N_2(t)\}$, $-\infty < t < \infty$, be a stationary bivariate point process that is strong mixing, $\alpha(\tau+u) = O\{\alpha(\tau)\}$ as $\tau \rightarrow \infty$, and such that $p_{ab}(u)$, $p_{abc}(u, v)$, $p_{abcd}(u, v, w)$ of (2.2) are finite and continuous for $a, b, c, d = 1, 2$. Then for $u_k^T \rightarrow u_k$, $|u_k^T - u_{k'}^T| \geq 2\beta$, $1 \leq k < k' \leq K$ and $\beta = L/T$, L constant, the variates $J_{a_1 b_1}^T(u_1^T), \dots, J_{a_K b_K}^T(u_K^T)$ are asymptotically independent $P\{2Lp_{a_k b_k}(u_k)\}$, $k = 1, \dots, K$ for $a_k, b_k = 1, 2$ as $T \rightarrow \infty$.

This result is proved in Section 4 of the paper assuming a direct variant of Theorem 1.3 of Volkonskii and Rozanov (1959). We may take $K = 1$ and $u_1^T = u$, here, and so see that $J_{ab}^T(u) \rightarrow P\{2Lp_{ab}(u)\}$ for $a, b = 1, 2$. We have allowed the arguments u_k^T to depend on T in order to be able to handle the case of a number of bins in the neighbourhood of a given lag u . The restriction on $|u_k^T - u_{k'}^T|$ means that the counting variates refer to distinct bins. In connection with the estimates of $p_{ab}(u)$, $m_{ab}(u)$ we have:

Corollary 1. Under the conditions of Theorem 1, $\hat{p}_{ab}(u)$, $\hat{p}'_{ab}(u)$, given by (1.7), (1.8), are asymptotically distributed as $(2L)^{-1}P\{2Lp_{ab}(u)\}$.

Corollary 2. Under the conditions of Theorem 1, $\hat{m}_{ab}(u)$, $\hat{m}'_{ab}(u)$, given by (1.7), (1.8), are asymptotically distributed as $(2L)^{-1}p_b^{-1}P\{2Lp_{ab}(u)\}$.

Had we so desired, we could have considered collections of estimates, at lags u_k^T , in the manner of the theorem, here. The asymptotic distributions of the estimates of (1.7) are not affected by the modification to (1.8) because of the convergence of the correction terms to zero, in probability. In both cases, the variance of the asymptotic distribution is seen to be proportional to the parameter being estimated. This occurrence suggests the application of a square root transformation to the estimates. We will return to this comment in the next section.

The estimates discussed here are histogram type estimates, involving a rectangular smoothing function. Cox (1965) remarks that one might want to consider other smoothing functions. For example, we might base estimates on

$$\sum_{i=-I}^I w_i J_{ab}^T(u - 2i\beta), \quad (2.3)$$

where $\sum w_i = 1$. From Theorem 1, the asymptotic distribution of this variate is seen to be that of $\sum w_i P_i$, where the P_i are independent $P\{2Lp_{ab}(u)\}$ variates. The mean of this asymptotic distribution is $2Lp_{ab}(u)$. The variance is $(\sum w_i^2) 2Lp_{ab}(u)$, a result that again suggests a square root transformation.

3. SOME FURTHER CONSIDERATIONS AND PRACTICAL EXAMPLES

The second-order product density, $p_{ab}(u)$, and the intensity function, $m_{ab}(u)$, both provide measures of the degree of statistical dependence of increments of the process $N_a(\cdot)$ that are u time units ahead of corresponding increments of the process $N_b(\cdot)$. In the case that these increments are independent, $p_{ab}(u) = p_a p_b$ and $m_{ab}(u) = p_a$. In the case that the process is strong mixing

$$E\{dN_a(t+u) dN_b(t)\} - E\{dN_a(t+u)\} E\{dN_b(t)\} = O\{\alpha(u)\} \rightarrow 0$$

as $|u| \rightarrow \infty$, see Volkonskii and Rozanov (1959) and so

$$\lim_{|u| \rightarrow \infty} p_{ab}(u) = p_a p_b \quad \text{and} \quad \lim_{|u| \rightarrow \infty} m_{ab}(u) = p_a. \quad (3.1)$$

This suggests that graphs of estimates of the functions $p_{ab}(\cdot)$ or $m_{ab}(\cdot)$ should also contain estimates of the constant levels $p_a p_b$ or p_a , as the case may be.

The relations of (3.1) suggest the source of the estimates $\hat{p}'_{ab}(u)$, $\hat{m}'_{ab}(u)$ of (1.8). For many processes, the covariance, $\text{cov}\{dN_a(t+u), dN_b(t)\}$ will be near 0 for large $|u|$. In the case of an ordinary bivariate stationary process $\{X_1(t), X_2(t)\}$ the covariance function

$$c_{ab}(u) = \text{cov}\{X_a(t+u), X_b(t)\}$$