

22S:138
Bayesian Statistics

Lecture 8
 Sept. 18, 2009

Kate Cowles
 374 SH, 335-0727
 kcowles@stat.uiowa.edu

Posterior predictive distribution of a future observation

- $p(y_{new}|y)$ is Normal
 - mean is posterior mean
 - variance is sum of
 - * data variance σ^2 (assumed known in this unrealistic case)
 - * variance of posterior mean

Jeffreys prior for normal mean with data variance assumed known

- $p(\mu) \propto 1, \quad -\infty < \mu < \infty$
- limit of $N(\mu_0, \sigma_0^2)$ as σ_0^2 goes to ∞
 σ_0^2 is prior variance
- equivalently, limit of $N(\mu_0, \tau_0^2)$ as τ_0^2 goes to 0
 τ_0^2 is prior precision

What is the posterior predictive distribution of the healing rate of a new newt?

Inference about the spread of a normal distribution

- primary research question may concern variability of response variable in population
 - quality control in industry
 - response to medical treatment

- Recall the joint sampling distribution of n observations modelled as conditionally independent draws from a normal

$$\begin{aligned}
 p(y_1, \dots, y_n | \mu, \sigma^2) &= \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma}} \exp\left[-\frac{(y_i - \mu)^2}{2\sigma^2}\right] \\
 &\propto \frac{1}{(\sigma^2)^{\frac{n}{2}}} \exp\left[-\frac{\sum_{i=1}^n (y_i - \mu)^2}{2\sigma^2}\right]
 \end{aligned}$$

- We will assume (unrealistically) that μ is a known constant

Inference for the variance of a normal distribution

- Suppose in the newt healing rate example that we knew the population mean $\mu = 25$ but we did not know the population variance σ^2
- Equivalently, we don't know the population precision τ^2
- We wish to infer about the distributions of these parameters that describe the spread of the normal distribution.

Sufficient statistic

- sufficient statistic for σ^2 is $\sum (y_i - \mu)^2$
- we can write likelihood equivalently as

$$p(y | \sigma^2) \propto \frac{1}{(\sigma^2)^{\frac{n}{2}}} \exp\left[-\frac{nv}{2\sigma^2}\right], \quad 0 < \sigma^2 < \infty$$

where

$$v = \frac{1}{n} \sum (y_i - \mu)^2$$

- What is corresponding conjugate prior?

Inverse gamma distribution

What is posterior?

$$p(\sigma^2|y) \propto ?$$

Estimating σ^2 of healing rates in population of newts

- Suppose μ was known to be 25.
- Suppose we had previously studied 2 newts and the average squared difference between their healing rate and 25 was 64.
- What is our appropriate prior?

• for newt data $\sum_{i=1}^{18} (y_i - 25)^2 = 1201$

- What is posterior

$$p(\sigma^2|y) \propto$$

- Recognize this as?

Alternative parameterization of prior

$$p(\sigma^2) = IG\left(\frac{\nu_0}{2}, \frac{\nu_0\sigma_0^2}{2}\right)$$

Then we can think of prior as providing equivalent information to

- ν_0 prior observations
- with σ_0^2 average squared deviation from known μ

Noninformative prior for normal variance

- What inverse gamma prior would have information equivalent to 0 prior observations?
- How would you write this as a p.d.f for σ^2 ?
- Is it proper or improper?
- What characteristics would a dataset have to have in order to produce a proper posterior distribution for σ^2 if this prior were used?